arXiv:1401.7843v1 [math-ph] 30 Jan 2014

Random Aharonov-Bohm vortices and some exact families of integrals: Part III

Stéphane Ouvry[1]

Université Paris-Sud, Laboratoire de Physique Théorique et Modèles Statistiques[2]
91405 Orsay, France

Abstract

As a sequel to [1] and [2], I present some recent progress on Bessel integrals $\int_0^\infty \mathrm{d}u\ uK_0(u)^n$, $\int_0^\infty \mathrm{d}u\ u^3K_0(u)^n$, ... where the power of the integration variable is odd and where n, the Bessel weight, is a positive integer. Some of these integrals for weights $n=3$ and $n=4$ are known to be intimately related to the zeta numbers $\zeta(2)$ and $\zeta(3)$. Starting from a Feynman diagram inspired representation in terms of n dimensional multiple integrals on an infinite domain, one shows how to partially integrate to $n-2$ dimensional multiple integrals on a finite domain. In this process the Bessel integrals are shown to be periods[3]. Interestingly enough, these "reduced" multiple integrals can be considered in parallel with some simple integral representations of ζ numbers. One also generalizes the construction of [2] on a particular sum of double nested Bessel integrals to a whole family of double nested integrals. Finally a strong PSLQ numerical evidence is shown to support a surprisingly simple expression of $\zeta(5)$ as a linear combination with rational coefficients of Bessel integrals of weight $n=8$.

1 Introduction: the quantum mechanics random magnetic impurity model and Bessel integrals

The random magnetic impurity model [3] describes a quantum particle in a plane coupled to a random distribution of Aharonov-Bohm fluxes perpendicular to the plane. It was introduced having in mind the Integer Quantum Hall effect. A perturbative expansion of the partition function of the model in the disorder coupling constant α (the Aharonov-Bohm flux expressed in unit of the quantum of flux) for 2 impurities, i.e. at second order in the impurity density ρ, lead us to consider Feynmann diagrams with maximal impurity line crossing at order $\rho^2\alpha^4$ [3] and $\rho^2\alpha^6$ [4].

On the one hand, these Feymann diagrams, which reduce after momenta integrations to multiple integrals on intermediate temperatures, were shown to rewrite in terms of simple integrals [5]

$$I_{\rho^2\alpha^4} = \int_0^\infty u\,K_0(u)^2(uK_1(u))^2\mathrm{d}u \tag{1}$$

[1] stephane.ouvry@u-psud.fr

[2] Unité Mixte de Recherche CNRS-Paris Sud, UMR 8626

[3] Periods are defined in [6] as "values of absolutely convergent integrals of rational functions with rational coefficients over domains in R^n given by polynomial inequalities with rational coefficients".

and double nested integrals [4]

$$\begin{aligned} I_{\rho^2\alpha^6} =& 8\int_0^\infty \mathrm{d}u\, u\, K_0(u)^2 (uK_1(u))^2 \int_0^u \mathrm{d}x\, xK_1(x)I_1(x)K_0(x)^2 \\ & - 4\int_0^\infty \mathrm{d}u\, uK_0(u)(uK_1(u))\,(uK_1(u)I_0(u) - uI_1(u)K_0(u)) \int_u^\infty \mathrm{d}x\, x\, K_0(x)^2 K_1(x)^2 \\ & + \int_0^\infty u\, K_0(u)^4 (uK_1(u))^2 \mathrm{d}u \end{aligned} \tag{2}$$

on products of modified Bessel functions K_ν and I_ν, with $\nu = 0, 1$.

On the other hand, one could show [5] by direct integration

$$\int_0^\infty uK_0(u)^4 \mathrm{d}u = \frac{2^3-1}{8}\zeta(3) \tag{3}$$

and by integration by part that

$$I_{\rho^2\alpha^4} = \int_0^\infty u\, K_0(u)^2 (uK_1(u))^2 \mathrm{d}u \tag{4}$$

is a linear combination with rational coefficients[4] of 1 and $\int_0^\infty uK_0(u)^4 \mathrm{d}u$, i.e. of 1 and $(2^3 - 1)\zeta(3)$. Similarly one obtained [4, 1] by direct integration[5]

$$\int_0^\infty \mathrm{d}u\, uI_0(u)K_0(u)^3 = \frac{2^2-1}{8}\zeta(2) \tag{5}$$

and by integration by part that

$$\int_0^\infty \mathrm{d}u\, uK_0(u)(uK_1(u))^2 I_0(u) \tag{6}$$

and

$$\int_0^\infty \mathrm{d}u\, uK_0(u)^2 uK_1(u)uI_1(u) \tag{7}$$

are linear combinations of 1 and $\int_0^\infty \mathrm{d}u\, uI_0(u)K_0(u)^3$, i.e. of 1 and $(2^2 - 1)\zeta(2)$. Likewise, by integration by part [2]

$$\int_0^\infty u\, K_0(u)^4 (uK_1(u))^2\, \mathrm{d}u = \frac{2}{15}\int_0^\infty u\, K_0(u)^6\, \mathrm{d}u - \frac{1}{5}\int_0^\infty u^3\, K_0(u)^6\, \mathrm{d}u \tag{8}$$

It was then natural to argue in [2] that the sum of double nested integrals (2) might also rewrite as a linear combination with rational coefficients of simple integrals on product of Bessel functions of weight 6 -defined as the total power of Bessel functions- and of ζ numbers of weigth $6 - 1 = 5$, like $\zeta(5)$, or below. The "counting rule" inferred from (3, 4, 5, 6, 7, 8) is that an integration $\int_0^\infty \mathrm{d}u\, u^n(\quad)$ with n odd diminishes the Bessel weight by one, so $\int_0^\infty u\, K_0(u)^4 (uK_1(u))^2$

[4] Linear combination with rational coefficients means here that there exist three positive or negative integers a, b and c such that

$$a\int_0^\infty u\, K_0(u)^2 (uK_1(u))^2 \mathrm{d}u + b\int_0^\infty uK_0(u)^4 \mathrm{d}u + c = 0$$

[5] See the Appendix for the derivations of (3) and (5).

is like $\zeta(5)$, that a Bessel function I_ν has no weight, so $\int_0^\infty \mathrm{d}u\, uI_0(u)K_0(u)^3$ is like $\zeta(2)$ and $\int_0^\infty \mathrm{d}u\, u^3\, K_0(u)^2K_1(u)^2 \int_0^u \mathrm{d}x\, xK_1(x)I_1(x)K_0(x)^2$ like $\zeta(3)\zeta(2)$ i.e. like $\zeta(5)$, that in turn is like $\int_0^\infty u\, K_0(u)^6$ or $\int_0^\infty u^3\, K_0(u)^6$ -for an explanation why only odd powers of u and no higher power appear here see [2] (see also subsection (2.3)). Indeed in [2] a numerical PSLQ [7] search gave

$$I_{\rho^2\alpha^6} =_{\mathrm{PSLQ}} \frac{1}{30}\int_0^\infty u\, K_0(u)^6\, \mathrm{d}u + \frac{1}{20}\int_0^\infty u^3 K_0(u)^6\, \mathrm{d}u - \frac{2^5-1}{160}\,\zeta(5) \tag{9}$$

(the 2^5-1 factor multiplying $\zeta(5)$ has to be viewed in parallel with 2^3-1 multiplying $\zeta(3)$ in (3) and 2^2-1 multiplying $\zeta(2)$ in (6)). Note that in (9), as in the sequel, an identity obtained from a numerical PSLQ search is labelled as $=_{\mathrm{PSLQ}}$.

It follows that the perturbative expansion of the random magnetic impurity model has Bessel integrals for building blocks. Note that Bessel integrals also appear in other physical contexts such as, for example, the Ising model [8].

In [2] it was argued that all integrals in the family

$$\int_0^\infty u^{p+1}K_0(u)^{n-j}(uK_1(u))^j\, \mathrm{d}u \qquad p \text{ even } \geq 0 \tag{10}$$

can be expressed as linear combinations with rational coefficients of a finite set of Bessel integrals $\int_0^\infty \mathrm{d}u\ uK_0(u)^n$, $\int_0^\infty \mathrm{d}u\ u^3K_0(u)^n$, ... , the latter constituting a basis for the former. In the present work one concentrates on different issues such as integrating the Bessel integrals.

2 Integrating $\int_0^\infty \mathrm{d}u\, uK_0(u)^n$

As alluded to in the Introduction, Feyman diagrams momenta integrations lead to multiple integrals on differences of consecutive intermediate temperatures. In the random magnetic impurity model, these multiple integrals take a form such that Bessel integrals can quite generally be representated in a similar way: change variable $u = 2\sqrt{t}$ and use the integral representation

$$\int_0^\infty \mathrm{d}a\, a^{\nu-1}e^{-a-\frac{t}{a}} = 2K_\nu(u)(\frac{u}{2})^\nu \tag{11}$$

so that

$$\begin{aligned}
&\int_0^\infty \mathrm{d}u\, uK_0(u)^n \\
=&\int_0^\infty 2\mathrm{d}t \int_0^\infty \frac{\mathrm{d}a_1}{2}\ldots\int_0^\infty \frac{\mathrm{d}a_n}{2}\frac{1}{a_1a_2\ldots a_n}e^{-a_1-\frac{t}{a_1}}e^{-a_2-\frac{t}{a_2}}\ldots e^{-a_n-\frac{t}{a_n}} \\
=&2\int_0^\infty \frac{\mathrm{d}a_1}{2}\ldots\int_0^\infty \frac{\mathrm{d}a_n}{2}\frac{1}{a_1a_2\ldots a_n}\frac{1}{\frac{1}{a_1}+\frac{1}{a_2}+\ldots+\frac{1}{a_n}}e^{-a_1-a_2-\ldots-a_n} \\
=&\frac{1}{2^{n-1}}\int_0^\infty \mathrm{d}a_1\ldots\int_0^\infty \mathrm{d}a_n\frac{1}{a_1a_2\ldots a_{n-1}+a_2a_3\ldots a_n+\ldots+a_na_1\ldots a_{n-2}}e^{-(a_1+a_2+\ldots+a_n)}
\end{aligned} \tag{12}$$

I am now going to show how to integrate the n dimensional multiple integral (12) to a $n-2$ multiple integral, which in turn will share some similarity with a well-known integral representation of ζ numbers. In the process the $n-1$ to $n-2$ variables reduction will be a non trivial example of a multidimensional integration leading to a simple and elegant expression. Note that it has so far not possible to integrate further along the same lines.

2.1 a_n integration

Introduce the variable β to rewrite $e^{-(a_1+a_2+\ldots+a_n)} = \int_0^\infty \mathrm{d}\beta\, e^{-\beta}\delta(\beta-(a_1+a_2+\ldots+a_n))$, then integrate over a_n

$$\begin{aligned}&\int_0^\infty \mathrm{d}u\, uK_0(u)^n\\ =&\frac{1}{2^{n-1}}\int_0^\infty \mathrm{d}\beta e^{-\beta}\int_0^\beta \mathrm{d}a_1\int_0^{\beta-a_1}\mathrm{d}a_2\ldots\int_0^{\beta-a_1-\ldots-a_{n-2}}\mathrm{d}a_{n-1}\\ &\frac{1}{a_1a_2\ldots a_{n-1}+(\beta-(a_1+a_2+\ldots+a_{n-1}))(a_2a_3\ldots a_{n-1}+a_3a_4\ldots a_{n-1}a_1+\ldots+a_1\ldots a_{n-2})}\end{aligned} \tag{13}$$

Change variables $a_i \to a_i' = a_i/\beta$ (notation $a_i' \to a_i$) so that the β integration becomes trivial and finally

$$\begin{aligned}&\int_0^\infty \mathrm{d}u\, uK_0(u)^n = \frac{1}{2^{n-1}}\int_0^1 \mathrm{d}a_1\int_0^{1-a_1}\mathrm{d}a_2\ldots\int_0^{1-a_1-\ldots-a_{n-2}}\mathrm{d}a_{n-1}\\ &\frac{1}{a_1a_2\ldots a_{n-1}+(1-(a_1+a_2+\ldots+a_{n-1}))(a_2a_3\ldots a_{n-1}+a_3a_4\ldots a_{n-1}a_1+\ldots+a_1a_2\ldots a_{n-2})}\end{aligned} \tag{14}$$

With the notations

$u_n = a_1+a_2+\ldots+a_n$

$v_n = a_2a_3\ldots a_n + a_3a_4\ldots a_na_1+\ldots+a_na_1a_2\ldots a_{n-2}+a_1a_2\ldots a_{n-1}$

$w_n = a_1a_2\ldots a_n$

one has shown that

$$\begin{aligned}\int_0^\infty \mathrm{d}u\, uK_0(u)^n &= \frac{1}{2^{n-1}}\int_0^\infty \mathrm{d}a_1\int_0^\infty \mathrm{d}a_2\ldots\int_0^\infty \mathrm{d}a_n\frac{1}{v_n}e^{-u_n}\\ &= \frac{1}{2^{n-1}}\int_0^1 \mathrm{d}a_1\int_0^{1-u_1}\mathrm{d}a_2\ldots\int_0^{1-u_{n-2}}\mathrm{d}a_{n-1}\frac{1}{w_{n-1}+(1-u_{n-1})v_{n-1}}\end{aligned} \tag{15}$$

2.2 a_{n-1} integration

Use

$$u_{n-1} = a_{n-1}+u_{n-2},\quad v_{n-1} = a_{n-1}v_{n-2}+w_{n-2},\quad w_{n-1} = a_{n-1}w_{n-2}$$

so that

$$\begin{aligned}&\frac{1}{w_{n-1}+(1-u_{n-1})v_{n-1}} = \frac{1}{w_{n-2}(1-u_{n-2})+a_{n-1}v_{n-2}(1-u_{n-2})-a_{n-1}^2v_{n-2}}\\ &= \frac{1}{-v_{n-2}(a_{n-1}-a_{n-1}^+)(a_{n-1}-a_{n-1}^-)} = \frac{1}{-v_{n-2}(a_{n-1}^+-a_{n-1}^-)}\left(\frac{1}{a_{n-1}-a_{n-1}^+}-\frac{1}{a_{n-1}-a_{n-1}^-}\right)\end{aligned} \tag{16}$$

where

$$a_{n-1}^\pm = \frac{-v_{n-2}(1-u_{n-2})\pm\sqrt{v_{n-2}^2(1-u_{n-2})^2+4w_{n-2}v_{n-2}(1-u_{n-2})}}{-2v_{n-2}}$$

One finds

$$\begin{aligned}&\int_0^\infty \mathrm{d}u\, uK_0(u)^n = \frac{1}{2^{n-1}}\int_0^1 \mathrm{d}a_1\int_0^{1-u_1}\mathrm{d}a_2\ldots\int_0^{1-u_{n-3}}\mathrm{d}a_{n-2}\\ &\frac{1}{\sqrt{v_{n-2}^2(1-u_{n-2})^2+4w_{n-2}v_{n-2}(1-u_{n-2})}}\log\left|\frac{a_{n-1}^-}{a_{n-1}^+}\frac{1-u_{n-2}-a_{n-1}^+}{1-u_{n-2}-a_{n-1}^-}\right|\end{aligned} \tag{17}$$

where the integrand rewrites as

$$\frac{2}{\sqrt{v_{n-2}^2(1-u_{n-2})^2+4w_{n-2}v_{n-2}(1-u_{n-2})}}\log|\frac{a_{n-1}^-}{a_{n-1}^+}|=\frac{2}{(1-u_{n-2})v_{n-2}}X\log\frac{1+X}{1-X} \tag{18}$$

with

$$X=\sqrt{\frac{1}{1+\frac{4w_{n-2}}{(1-u_{n-2})v_{n-2}}}} \tag{19}$$

At this point a seemingly complicated -square root- expression of the remaining $n-2$ variables appears. However the change of variables

$$a_i\to x_i=X\frac{a_i}{u_{n-2}} \tag{20}$$

so that when $0<a_i<1$ then $1>X>0\Rightarrow 1>x_i>0$ will give a more tractable form for the integrand. The Jacobian is

$$\frac{(u_{n-2})^{n-2}}{X^{n-3}|\sum_{i=1}^{n-2}a_i\partial_i X|}$$

Use homogeneity relations

$$\begin{aligned}\sum_{i=1}^{n-2}a_i\partial_i u_{n-2}&=u_{n-2}\\ \sum_{i=1}^{n-2}a_i\partial_i v_{n-2}&=(n-3)v_{n-2}\\ \sum_{i=1}^{n-2}a_i\partial_i w_{n-2}&=(n-2)w_{n-2}\end{aligned} \tag{21}$$

so that

$$\sum_{i=1}^{n-2}a_i\partial_i X=-\frac{1}{2}X(1-X^2)\frac{1}{1-u_{n-2}}$$

and

$$\begin{aligned}\int_0^\infty \mathrm{d}u\, uK_0(u)^n=&\frac{1}{2^{n-1}}\int_0^1\mathrm{d}x_1\int_0^{1-x_1}\mathrm{d}x_2\ldots\int_0^{1-x_1-\ldots-x_{n-3}}\mathrm{d}x_{n-2}\log\frac{1+X}{1-X}\\ &\frac{2}{(1-u_{n-2})v_{n-2}}X\frac{(u_{n-2})^{n-2}}{X^{n-3}|-\frac{1}{2}X(1-X^2)\frac{1}{1-u_{n-2}}|}\\ =&\frac{1}{2^{n-1}}\int_0^1\mathrm{d}x_1\int_0^{1-x_1}\mathrm{d}x_2\ldots\int_0^{1-x_1-\ldots-x_{n-3}}\mathrm{d}x_{n-2}\log\frac{1+X}{1-X}\\ &\frac{4}{v_{n-2}}\frac{(u_{n-2})^{n-2}}{X^{n-3}(1-X^2)}\end{aligned} \tag{22}$$

From (19) and (20) it follows

$$\begin{aligned}X&=x_1+x_2+\ldots+x_{n-2}\\ w_{n-2}X^{n-2}&=(u_{n-2})^{n-2}x_1x_2\ldots x_{n-2}\\ \frac{u_{n-2}v_{n-2}}{w_{n-2}X}&=\frac{1}{x_1}+\frac{1}{x_2}+\ldots+\frac{1}{x_{n-2}}\\ \frac{1}{X^2}-1&=\frac{4w_{n-2}}{(1-u_{n-2})v_{n-2}}\end{aligned} \tag{23}$$

so that finally $\int_0^\infty \mathrm{d}u\, uK_0(u)^n$ takes the desired simple form

$$\int_0^\infty \mathrm{d}u\, uK_0(u)^n = \frac{1}{2^{n-1}} \int_0^1 \mathrm{d}x_1 \int_0^{1-x_1} \mathrm{d}x_2 \ldots \int_0^{1-x_1-\ldots-x_{n-3}} \mathrm{d}x_{n-2} \log \frac{1+x_1+x_2+\ldots+x_{n-2}}{1-(x_1+x_2+\ldots+x_{n-2})} \frac{4}{4(x_1+\ldots+x_{n-2})x_1\ldots x_{n-2}+(1-(x_1+\ldots+x_{n-2})^2)(x_2x_3\ldots x_{n-2}+\ldots+x_1x_2\ldots x_{n-3})} \tag{24}$$

This procedure generalises to $\int_0^\infty \mathrm{d}u\, u^p K_0(u)^n$ with p odd: for example for $p=3$ one obtains similarly

$$\int_0^\infty \mathrm{d}u\, u^3 K_0(u)^n = \frac{1}{2^{n-3}} \int_0^1 \mathrm{d}a_1 \int_0^{1-a_1} \mathrm{d}a_2 \ldots \int_0^{1-a_1-\ldots-a_{n-2}} \mathrm{d}a_{n-1} \frac{a_1a_2\ldots a_{n-1}(1-(a_1+a_2+\ldots+a_{n-1}))}{(a_1a_2\ldots a_{n-1}+(1-(a_1+a_2+\ldots+a_{n-1}))(a_2a_3\ldots a_{n-1}+a_3a_4\ldots a_{n-1}a_1+\ldots+a_1a_2\ldots a_{n-2}))^2}$$
$$= \frac{1}{2^{n-3}} \int_0^1 \mathrm{d}x_1 \int_0^{1-x_1} \mathrm{d}x_2 \ldots \int_0^{1-x_1-\ldots-x_{n-3}} \mathrm{d}x_{n-2} \left(\frac{1+(x_1+x_2+\ldots+x_{n-2})^2}{2(x_1+x_2+\ldots+x_{n-2})} \log \frac{1+x_1+x_2+\ldots+x_{n-2}}{1-(x_1+x_2+\ldots+x_{n-2})} - 1 \right) \frac{4x_1\ldots x_{n-2}\left(1-(x_1+x_2+\ldots+x_{n-2})^2\right)}{\left(4(x_1+\ldots+x_{n-2})x_1\ldots x_{n-2}+(1-(x_1+\ldots+x_{n-2})^2)(x_2x_3\ldots x_{n-2}+\ldots+x_1x_2\ldots x_{n-3})\right)^2} \tag{25}$$

2.3 Some remarks

(24) and (25) indicate that $\int_0^\infty \mathrm{d}u\, uK_0(u)^n$ and $\int_0^\infty \mathrm{d}u\, u^3K_0(u)^n$ are periods. This is the case in general for $\int_0^\infty \mathrm{d}u\, u^pK_0(u)^n$ with p odd. Note that it is crucial the power p of the integration variable u be odd to obtain rational functions of the $n-2$ remaining variables $x_1,\ldots,x_{n-2}$ -if not square root would appear in the process.

Looking at (24), it has been found that $\int_0^\infty \mathrm{d}u\, uK_0(u)^n$ rewrites as a $n-2$ integral on the finite domain $0 \leq x_1+x_2+\ldots x_{n-2} \leq 1$ with, for integrand, $\log \frac{1+x_1+x_2+\ldots+x_{n-2}}{1-(x_1+x_2+\ldots+x_{n-2})}$ multiplied by a rational function. Interestingly enough, this structure shares some similarities with the well-known rewriting of ζ numbers in terms of a simple integral on the domain $0 \leq x \leq 1$ with for integrand $(\log \frac{1+x}{1-x})^{n-1}$ multiplied by $1/x$

$$\frac{2^n-1}{2^{n-1}}\zeta(n) = \sum_{k=1}^\infty \frac{1}{k^n} + (-1)^{k-1}\frac{1}{k^n} = \frac{1}{2}\int_0^\infty \mathrm{d}x x^{n-1}\left(\frac{1}{e^x-1}+\frac{1}{e^x+1}\right) = \frac{1}{(n-1)!}\int_0^1 \mathrm{d}x (\log\frac{1+x}{1-x})^{n-1}\frac{1}{x} \tag{26}$$

One can argue in (24) that the logarithm has been linearized in place of appearing in (26) at a power $n-1$, but the price to be paid is a multiple integral on $n-2$ variables with the variable x in (26) replaced in the logarithm by the sum $x_1+x_2+\ldots+x_{n-2}$ and in the factor $1/x$ by a more involved polynomial of degree $n-1$.

In the simple cases $n=3$ and $n=4$, one can indeed integrate (24) completely to simple expressions in terms of $\zeta(2)$ and $\zeta(3)$ numbers (see (3) and (44, 48) in the Appendix).

2.4 A summary

Finally if in (24) and (25) one changes notations $x_i \to a_i$ one has shown

$$2^{n-1}\int_0^\infty \mathrm{d}u\; uK_0(u)^n = \int_0^\infty \mathrm{d}a_1 \ldots \int_0^\infty \mathrm{d}a_{n-2}\int_0^\infty \mathrm{d}a_{n-1}\int_0^\infty \mathrm{d}a_n \quad \frac{1}{v_n} \quad e^{-u_n}$$
$$= \int_0^1 \mathrm{d}a_1 \int_0^{1-u_1} \mathrm{d}a_2 \ldots \int_0^{1-u_{n-3}} \mathrm{d}a_{n-2}\int_0^{1-u_{n-2}} \mathrm{d}a_{n-1} \quad \frac{1}{w_{n-1}+(1-u_{n-1})v_{n-1}} \tag{27}$$
$$= \int_0^1 \mathrm{d}a_1 \int_0^{1-u_1} \mathrm{d}a_2 \ldots \int_0^{1-u_{n-3}} \mathrm{d}a_{n-2} \quad \log\frac{1+u_{n-2}}{1-u_{n-2}} \quad \frac{4}{4u_{n-2}w_{n-2}+(1-u_{n-2}^2)v_{n-2}} \tag{28}$$

and

$$2^{n-3}\int_0^\infty \mathrm{d}u\; u^3K_0(u)^n = \int_0^\infty \mathrm{d}a_1 \ldots \int_0^\infty \mathrm{d}a_{n-2}\int_0^\infty \mathrm{d}a_{n-1}\int_0^\infty \mathrm{d}a_n \quad \frac{w_n}{v_n^2} \quad e^{-u_n}$$
$$= \int_0^1 \mathrm{d}a_1 \int_0^{1-u_1} \mathrm{d}a_2 \ldots \int_0^{1-u_{n-3}} \mathrm{d}a_{n-2}\int_0^{1-u_{n-2}} \mathrm{d}a_{n-1} \quad \frac{w_{n-1}(1-u_{n-1})}{(w_{n-1}+(1-u_{n-1})v_{n-1})^2} \tag{29}$$
$$= \int_0^1 \mathrm{d}a_1 \int_0^{1-u_1} \mathrm{d}a_2 \ldots \int_0^{1-u_{n-3}} \mathrm{d}a_{n-2} \left(\frac{1+u_{n-2}^2}{2u_{n-2}}\log\frac{1+u_{n-2}}{1-u_{n-2}} - 1\right) \frac{4w_{n-2}(1-u_{n-2}^2)}{(4u_{n-2}w_{n-2}+(1-u_{n-2}^2)v_{n-2})^2}$$

As a remark one can proceed similarly with $\int_0^\infty \mathrm{d}u\, uI_0(u)K_0(u)^n$: using again the integral representation (11) and ($u = 2\sqrt{t}$)

$$I_0(u) = \sum_{k=0}^\infty \frac{t^k}{(k!)^2} \tag{30}$$

one can integrate over t using

$$\int_0^\infty \mathrm{d}t\; t^k e^{-tx} = \frac{k!}{x^{k+1}} \tag{31}$$

to obtain

$$2^{n-1}\int_0^\infty \mathrm{d}u\; uI_0(u)K_0(u)^n = \int_0^\infty \mathrm{d}a_1 \ldots \int_0^\infty \mathrm{d}a_{n-2}\int_0^\infty \mathrm{d}a_{n-1}\int_0^\infty \mathrm{d}a_n \quad \frac{1}{v_n} \quad e^{-u_n+\frac{w_n}{v_n}}$$

Introducing as above the variable β, integrating over a_n and then trivially over β one finally obtains

$$2^{n-1}\int_0^\infty \mathrm{d}u\; uI_0(u)K_0(u)^n$$
$$= \int_0^1 \mathrm{d}a_1 \int_0^{1-u_1} \mathrm{d}a_2 \ldots \int_0^{1-u_{n-3}} \mathrm{d}a_{n-2}\int_0^{1-u_{n-2}} \mathrm{d}a_{n-1} \quad \frac{1}{w_{n-1}u_{n-1}+(1-u_{n-1})v_{n-1}} \tag{32}$$

a result to be compared to (15). One could push the integration a step further to obtain an expression in terms of u_{n-2}, v_{n-2} and w_{n-2} but somehow more involved than (28) since its denominator would contain a square root.

3 Rewriting $I_{\rho^2\alpha^6}$ and a family of double nested Bessel integrals

The fact that the double nested integrals in (2) can be reexpressed in (9) as a linear combination with rational coefficients of simple Bessel integrals with the right weight -here weight 6- and of $\zeta(5)$, fits well in the "Bessel integral $\to$ zeta number" mapping [2]. If a product f of Bessel functions is, like in (3, 4, 5, 6, 7), mapped by simple integration on a ζ number denoted by $\tilde{\zeta}(f)$

$$f \to \int_0^\infty f(u)\mathrm{d}u = \tilde{\zeta}(f)$$

then for a pair of such products f, g the mapping by double "nested" integration

$$f, g \to \int_0^\infty f(u)\mathrm{d}u \int_0^u g(x)\mathrm{d}x = \tilde{\zeta}(f, g)$$

on a polyzeta number denoted by $\tilde{\zeta}(f, g)$ makes sense, since, because of

$$\int_0^\infty f(u)\mathrm{d}u \int_0^u g(x)\mathrm{d}x = \int_0^\infty f(u)\mathrm{d}u \int_0^\infty g(x)\mathrm{d}x - \int_0^\infty g(u)\mathrm{d}u \int_0^u f(x)\mathrm{d}x,$$

one has

$$\tilde{\zeta}(f, g) = \tilde{\zeta}(f)\tilde{\zeta}(g) - \tilde{\zeta}(g, f) \tag{33}$$

in analogy with

$$\zeta(p, q) = \zeta(p)\zeta(q) - \zeta(p + q) - \zeta(q, p) \tag{34}$$

for the standard polyzeta $\zeta(p, q) = \sum_{n>m} \frac{1}{n^p}\frac{1}{m^q}$ - if $\zeta(p, q)$ would be defined as $\sum_{n>m} \frac{1}{n^p}\frac{1}{m^q} + \frac{1}{2}\zeta(p+q)$ then (34) would take the form (33).

Coming back to (2,9) use

$$\int_0^\infty g(u)\mathrm{d}u \int_u^\infty f(x)\mathrm{d}x = \int_0^\infty f(u)\mathrm{d}u \int_0^u g(x)\mathrm{d}x$$

to rewrite $I_{\rho^2\alpha^6}$ as

$$\begin{aligned} I_{\rho^2\alpha^6} = 8 &\int_0^\infty \mathrm{d}u\, u\, K_0(u)^2 (uK_1(u))^2 \int_0^u \mathrm{d}x\, xK_1(x)I_1(x)K_0(x)^2 \\ -4 &\int_0^\infty \mathrm{d}u\, u\, K_0(u)^2 K_1(u)^2 \int_0^u \mathrm{d}x\, xK_0(x)xK_1(x)\big(xK_1(x)I_0(x) - xK_0(x)I_1(x)\big) \\ + &\int_0^\infty u\, K_0(u)^4 (uK_1(u))^2 \mathrm{d}u \end{aligned} \tag{35}$$

Next use $xI_1(x)K_0(x) + xI_0(x)K_1(x) = 1$ so that

$$\begin{aligned} I_{\rho^2\alpha^6} = 8 &\int_0^\infty \mathrm{d}u\, u^3\, K_0(u)^2 K_1(u)^2 \int_0^u \mathrm{d}x\, xK_1(x)I_1(x)K_0(x)^2 \\ +8 &\int_0^\infty \mathrm{d}u\, u\, K_0(u)^2 K_1(u)^2 \int_0^u \mathrm{d}x\, x^3 K_1(x)I_1(x)K_0(x)^2 \\ -4 &\int_0^\infty \mathrm{d}u\, u\, K_0(u)^2 K_1(u)^2 \int_0^u \mathrm{d}x\, x^2 K_0(x)K_1(x) \\ + &\int_0^\infty u\, K_0(u)^4 (uK_1(u))^2 \mathrm{d}u \end{aligned} \tag{36}$$

One has for the next to last term in (36)

$$\begin{aligned} -4 \int_0^\infty \mathrm{d}u\, u\, K_0(u)^2 K_1(u)^2 \int_0^u \mathrm{d}x\, x^2 K_0(x)K_1(x) &= 2 \int_0^\infty u\, K_0(u)^2 K_1(u)^2 \big((uK_1(u))^2 - 1\big)\mathrm{d}u \\ &= -\frac{2}{3} \int_0^\infty u\, K_1(u) \big((uK_1(u))^2 - 1\big)\mathrm{d}K_0(u)^3 \\ &= \frac{2}{3} \int_0^\infty K_0(u)^3\, \mathrm{d}(uK_1(u))\big((uK_1(u))^2 - 1\big) \\ &= \frac{2}{3} \int_0^\infty uK_0(u)^4 \mathrm{d}u\, - 2 \int_0^\infty u\, K_0(u)^4 (uK_1(u))^2 \mathrm{d}u \end{aligned} \tag{37}$$

so that

$$\begin{aligned} I_{\rho^2\alpha^6} &= 8\int_0^\infty \mathrm{d}u\, u^3\, K_0(u)^2 K_1(u)^2 \int_0^u \mathrm{d}x\, x K_1(x) I_1(x) K_0(x)^2 \\ &+8\int_0^\infty \mathrm{d}u\, u\, K_0(u)^2 K_1(u)^2 \int_0^u \mathrm{d}x\, x^3 K_1(x) I_1(x) K_0(x)^2 \\ &+\frac{2}{3}\int_0^\infty u K_0(u)^4 \mathrm{d}u \\ &-\int_0^\infty u\, K_0(u)^4 (u K_1(u))^2 \mathrm{d}u \end{aligned} \tag{38}$$

Defining

$$f_n(x) = x^n K_0(x)^2 K_1(x)^2 \quad \text{and} \quad g_n(x) = x^n K_1(x) I_1(x) K_0(x)^2$$

equations (4, 8, 38) imply that

$$\begin{aligned} \tilde{\zeta}(f_3, g_1) + \tilde{\zeta}(f_1, g_3) =& \frac{1}{48}\int_0^\infty u\, K_0(u)^6\, \mathrm{d}u - \frac{3}{160}\int_0^\infty u^3 K_0(u)^6\, \mathrm{d}u \\ &- \frac{7}{96}\zeta(3) - \frac{31}{1280}\zeta(5) \end{aligned} \tag{39}$$

as confirmed by a direct PSLQ check.

So the meaning of (9) when rewritten as (39) might be that it is the symmetric form $\tilde{\zeta}(f_3, g_1)+\tilde{\zeta}(f_1, g_3)$ which is a linear combination with rational coefficients of simple Bessel integrals of weigth 6 and ζ numbers like $\zeta(5)$ and below. This suggests to look at other symmetric sums of double nested integrals sharing this property. A PSLQ search confirms that $\tilde{\zeta}(f_5, g_1) + \tilde{\zeta}(f_1, g_5)$ belongs indeed to this category

$$\begin{aligned} \tilde{\zeta}(f_5, g_1) + \tilde{\zeta}(f_1, g_5) =_{PSLQ}& \frac{211}{11520}\int_0^\infty u\, K_0(u)^6\, \mathrm{d}u + \frac{3953}{23040}\int_0^\infty u^3 K_0(u)^6\, \mathrm{d}u \\ &+ \frac{11}{9216} - \frac{1}{9}\zeta(3) - \frac{93}{5120}\zeta(5) \end{aligned} \tag{40}$$

as well as, for example, $\tilde{\zeta}(f_7, g_1) + \tilde{\zeta}(f_1, g_7)$

$$\begin{aligned} \tilde{\zeta}(f_7, g_1) + \tilde{\zeta}(f_1, g_7) =_{PSLQ}& \frac{108731}{1728000}\int_0^\infty u\, K_0(u)^6\, \mathrm{d}u + \frac{4256617}{3456000}\int_0^\infty u^3 K_0(u)^6\, \mathrm{d}u \\ &+ \frac{27877}{460800} - \frac{8}{15}\zeta(3) - \frac{279}{5120}\zeta(5) \end{aligned} \tag{41}$$

and $\tilde{\zeta}(f_3, g_5) + \tilde{\zeta}(f_5, g_3)$

$$\begin{aligned} \tilde{\zeta}(f_3, g_5) + \tilde{\zeta}(f_5, g_3) =_{PSLQ}& -\frac{28921}{691200}\int_0^\infty u\, K_0(u)^6\, \mathrm{d}u + \frac{1151533}{1382400}\int_0^\infty u^3 K_0(u)^6\, \mathrm{d}u \\ &+ \frac{14653}{184320} + \frac{25}{192}\zeta(3) + \frac{279}{20480}\zeta(5) \end{aligned} \tag{42}$$

One infers that for any two odd positive integers n, m one should have that $\tilde{\zeta}(f_n, g_m) + \tilde{\zeta}(f_m, g_n)$ can be rewritten as a linear combination of 1, $\int_0^\infty u\, K_0(u)^6\, \mathrm{d}u$, $\int_0^\infty u^3\, K_0(u)^6\, \mathrm{d}u$, $\zeta(3)$ and $\zeta(5)$. Finding the coefficients of the linear combination can be in principle achieved by generalizing the integration by part procedures [1, 2, 3, 5] for simple integrals to double nested integrals.

4 Conclusion : Why looking at Bessel integrals?

In Section (1) some arguments (and examples) were given of why Bessel integrals are expected to be related to ζ numbers.

On the one hand, one has found the surprisingly simple PSLQ identity

$$\zeta(5) =_{\rm PSLQ} \frac{1}{77}\int_0^\infty u\,K_0(u)^8\,{\rm d}u - \frac{72}{77}\int_0^\infty u^3\,K_0(u)^8\,{\rm d}u \tag{43}$$

It would certainly be rewarding to look for a sytematics, if any, behind, for example, (3), (43) and (44).

On the other hand, one has tried, to this aim, to integrate the Bessel integrals starting from Feynman diagram inspired multidimensional integrals on an infinite domain. In the process on has been able to reduce these integrals via a non trivial multidimensional integration to $n-2$ dimensional integrals on a finite domain, whose structure are reminiscent of well-known simple integral representations of ζ numbers.

One would certainly like to see at integrating $\int_0^\infty u\,K_0(u)^n\,{\rm d}u$ further up to possibly one integration left on an integrand which might contain, as in (26), a term like $(\log\frac{1+x}{1-x})^{n-1}$ times a rational function of x yet to be determined, possibly allowing for a derivation of (43), and, more generally, of relations between Bessel integrals and ζ (and polyzeta) numbers.

Finally, the fact that Bessel integrals fall in the category of periods might also be an indication of a deeper meaning yet to be understood.

Acknowledgments: I acknowledge some useful conversations with Stefan Mashkevich in particular for helping in the numerics involved in the PSLQ searches of Section (3). I have also benefited from numerous technical and helpful discussions with Jean Desbois.

Appendix : More on integrating

n = 3

From (24)

$$\begin{aligned}\int_0^\infty \mathrm{d}u\, uK_0(u)^3 &= \int_0^1 \mathrm{d}x_1 \log\frac{1+x_1}{1-x_1}\frac{1}{1+3x_1^2} = \int_0^1 \mathrm{d}x_1 \log\frac{1+x_1}{1-x_1}\frac{2}{(1+x_1)^3(1+(\frac{1-x_1}{1+x_1})^3)}\\ &= -\frac{1}{2}\int_0^1 \mathrm{d}u \frac{1+u}{1+u^3}\log u\\ &= -\frac{1}{2}\int_0^1 \mathrm{d}u(1+u)\sum_{k=0}^\infty (-1)^k u^{3k}\log u\\ &= \frac{1}{2}\sum_{k=0}^\infty (-1)^k\left(\frac{1}{(3k+1)^2}+\frac{1}{(3k+2)^2}\right) = \frac{\psi_1(1/3)-\psi_1(2/3)}{12}\end{aligned} \tag{44}$$

where one has made the change of variable $u = \frac{1-x_1}{1+x_1}$ and used

$$\int_0^1 \mathrm{d}u\, u^k(\log u)^n = (-1)^n\frac{n!}{(k+1)^n} \tag{45}$$

In (44) $\psi_1(z)$ stands for the polygamma function such that $8\zeta(2) = \psi_1(1/3)+\psi_1(2/3)$, thus relating $\int_0^\infty \mathrm{d}u\, uK_0(u)^3$ to $\zeta(2)$.

From(32)

$$\int_0^\infty \mathrm{d}u\, uI_0(u)K_0(u)^3 = \frac{1}{4}\int_0^1 \mathrm{d}x_1\int_0^{1-x_1}\mathrm{d}x_2 \frac{1}{x_1x_2(x_1+x_2)-(x_1+x_2)(1-x_1-x_2)} \tag{46}$$

With the change of variables $x = x_1+x_2$ and $y = x_1-x_2$ one gets

$$\begin{aligned}\int_0^\infty \mathrm{d}u\, uI_0(u)K_0(u)^3 =& \frac{1}{4}\int_0^1 \mathrm{d}x\int_0^x \mathrm{d}y \frac{4}{x}\frac{1}{x-2+y}\frac{1}{x-2-y}\\ =& \frac{1}{2}\int_0^1 \mathrm{d}x\frac{1}{x(x-2)}\int_0^x \mathrm{d}y\frac{1}{x-2+y}+\frac{1}{x-2-y}\\ =& \frac{1}{2}\int_0^1 \mathrm{d}x\frac{1}{x(x-2)}\log(1-x)\\ =& -\frac{1}{2}\int_0^1 \mathrm{d}u\frac{1}{1-u^2}\log u = \frac{3\zeta(2)}{8}\end{aligned} \tag{47}$$

where one has set $u = 1-x$.

n = 4

From (24)

$$\begin{aligned}\int_0^\infty \mathrm{d}u\, uK_0(u)^4 &= \frac{1}{2^3}\int_0^1 \mathrm{d}x_1 \int_0^{1-x_1} \mathrm{d}x_2 \log\frac{1+x_1+x_2}{1-x_1-x_2}\frac{1}{x_1+x_2}\frac{4}{1-(x_1+x_2)^2+4x_1x_2}\\ &= \frac{1}{2^3}\frac{1}{2}\int_0^1 \mathrm{d}x\frac{1}{x}\log\frac{1+x}{1-x}\int_{-x}^{x}\mathrm{d}y\frac{4}{1-y^2}\\ &= \frac{1}{2}\int_0^1 \mathrm{d}x\frac{1}{x}\log\frac{1+x}{1-x}\int_0^{x}\mathrm{d}y\frac{1}{1-y^2}\\ &= \frac{1}{2}\frac{1}{2}\int_0^1 \mathrm{d}x\frac{1}{x}(\log\frac{1+x}{1-x})^2\\ &= \frac{1}{2}\int_0^1 \mathrm{d}u\frac{1}{1-u^2}(\log u)^2\\ &= \frac{1}{2}\int_0^1 \mathrm{d}u(\log u)^2\sum_{n=0}^{\infty}u^{2n} = \frac{7\zeta(3)}{8}\end{aligned} \tag{48}$$

with the change of variable $u=\frac{1-x}{1+x}$.

As a remark one knows (see [1]) that $\int_0^\infty \mathrm{d}u\, u^3K_0(u)^4$ is a linear combination with rational coefficients of 1 and $\int_0^\infty \mathrm{d}u\, uK_0(u)^4$ namely

$$4\int_0^\infty \mathrm{d}u\, uK_0(u)^4 - 16\int_0^\infty \mathrm{d}u\, u^3K_0(u)^4 = 3 \tag{49}$$

From (25)

$$\begin{aligned}\int_0^\infty \mathrm{d}u\, u^3K_0(u)^4 = \int_0^1 \mathrm{d}x_1 \int_0^{1-x_1} \mathrm{d}x_2 \left(\frac{1+(x_1+x_2)^2}{2(x_1+x_2)}\log\frac{1+x_1+x_2}{1-(x_1+x_2)}-1\right)\frac{1-(x_1+x_2)^2}{(x_1+x_2)^2}\\ \frac{4x_1x_2}{\left(1-(x_1+x_2)^2+4x_1x_2\right)^2}\end{aligned} \tag{50}$$

so that (49) implies that

$$\begin{aligned}&\int_0^1 \mathrm{d}x_1 \int_0^{1-x_1} \mathrm{d}x_2 \log\frac{1+x_1+x_2}{1-x_1-x_2}\frac{1}{x_1+x_2}\frac{2}{1-(x_1+x_2)^2+4x_1x_2}\\ &-8\left(\frac{1+(x_1+x_2)^2}{2(x_1+x_2)}\log\frac{1+x_1+x_2}{1-(x_1+x_2)}-1\right)\frac{1-(x_1+x_2)^2}{(x_1+x_2)^2}\frac{4x_1x_2}{\left(1-(x_1+x_2)^2+4x_1x_2\right)^2} = 3\end{aligned} \tag{51}$$

has to be satisfied.

With the change of variables $x=x_1+x_2$ and $y=x_1-x_2$ the right hand side of (50) becomes

$$\begin{aligned}&\int_0^1 \mathrm{d}x\left(\frac{1+x^2}{2x}\log\frac{1+x}{1-x}-1\right)\frac{1-x^2}{x^2}\frac{1}{2}\left(-x+(1+x^2)\frac{1}{2}\log\frac{1+x}{1-x}\right)\\ &= \int_0^1 \mathrm{d}x\frac{1-x^2}{2x}\left(\frac{1+x^2}{2x}\log\frac{1+x}{1-x}-1\right)^2\end{aligned} \tag{52}$$

Finally (51), becomes

$$\int_0^1 \mathrm{d}x\left(\frac{1}{x}(\log\frac{1+x}{1-x})^2 - 4\frac{1-x^2}{x}(\frac{1+x^2}{2x}\log\frac{1+x}{1-x}-1)^2\right) = 3 \tag{53}$$

which is indeed true.

References

[1] S. Ouvry, Random Aharonov-Bohm vortices and some exact families of integrals, JSTAT: Theory and Experiment (2005) P09004.

[2] S. Mashkevich and S. Ouvry, Random Aharonov-Bohm vortices and some exact families of integrals: Part II, JSTAT: Theory and Experiment (2008) P03018.

[3] J. Desbois, C. Furtlehner and S. Ouvry, Random Magnetic Impurities and the Landau Problem, Nuclear Physics B[FS] **453** (1995) 759.

[4] C. Furtlehner and S. Ouvry, private communication.

[5] C. Furtlehner and S. Ouvry, Integrals involving four Macdonald functions and their relation to $7\zeta(3)/2$, arXiv:math-ph/0306004.

[6] M. Konsevitch and D. Zagier, Periods (2001) IHES/M/01/22.

[7] http://mathworld.wolfram.com/PSLQAlgorithm.html, and references therein.

[8] David H. Bailey, Jonathan M. Borwein, David Broadhurst, M. L. Glasser, Elliptic integral evaluations of Bessel moments, arXiv:0801.0891. J. Phys. A: Math. Theor. 41 (2008) 205203.